

- Arboric. 3:114-118.
11. Shortle, W.C. 1984. *Biochemical mechanisms of discoloration, decay, and compartmentalisation of decay in trees*. Inter. Assoc. Wood Anat. Bull. n.s. 5:100-104.
  12. Smith, K.T., R.O. Blanchard, and W.C. Shortle. 1981. *Postulated mechanism of biological control of decay fungi in red maple wounds treated with Trichoderma harzianum*. Phytopathology 71:496-498.
  13. Smith, K.T. and W.C. Shortle. 1988. *Electrical*

*resistance and wood decay by white rot fungi*. Mycologia 80:124-126.

USDA Forest Service  
 Northeastern Forest Experiment Station  
 P.O. Box 640  
 Durham, New Hampshire 03824

## THERAPY FOR DUTCH ELM DISEASE<sup>1</sup>

by Gerald N. Lanier

**Abstract.** An aggregate of 82 American elms in Syracuse New York and Washington D.C. naturally infected with Dutch elm disease were given therapy by pruning infected limbs, injection of benzimidazol fungicides, or combinations of these treatments. Pruning alone was applied only when there was a distance of 3 m or more from the last visible streak to the distal cut. Pruning without injection was successful in each of 10 current year infections, but in none of 3 residual infections. Fungicide injection without pruning succeeded in 76% of current year and 33% of residual infections. Fungicide injection plus pruning was successful therapy for 100% of the current year and 71% of the residual infections. Each of five "incurable" trees showed no further symptoms after fungicide was injected directly into infected wood in a large limb or the bole as well as into the root collars. Relatively massive injection of fungicide in trees with root graft-transmitted DED did not succeed.

**Résumé.** Un ensemble de 82 ormes américains à Syracuse, New York et à Washington, D. C., infectées

naturellement par la maladie hollandaise de l'orme, reçurent un traitement consistant en un élagage des branches infectées, une injection de fongicides à base de benzimidazol, ou une combinaison de ces traitements. L'élagage seul fut appliqué seulement lorsque qu'il y avait une distance de 3 m ou plus de la dernière strie visible à la coupe. L'élagage sans injection fut un succès pour chacune des 10 infections de l'année courante, mais pour aucune des 3 infections résiduelles. L'injection d'un fongicide sans élagage a réussi dans 76% des infections de l'année en cours et dans 33% des infections résiduelles. L'injection d'un fongicide avec un élagage fut un traitement efficace à 100% pour les infections de l'année en cours et à 71% pour les infections résiduelles. Cinq arbres "incurable" n'ont pas présenté d'autres symptômes après que le fongicide fut injecté dans le bois infecté dans une grosse branche ou dans de tronc, de même qu'à la base de l'arbre. Une injection massive de fongicides dans des arbres infectés par des greffes de racines n'a pas réussi.

1. Presented at the Symposium on Systemic Chemical Treatments in Tree Culture at Michigan State University, East Lansing in October 1987.

Therapy of elms affected with Dutch elm disease (DED) by eradicated pruning (1, 2, 6) or injection of benzimidazole fungicides (7, 8, 9, 10) is well established. By either method, rates of success have generally been predicted by the percent of the tree crown showing DED symptoms at the time of treatment. Guidelines for injection generally state trees that have less than 5-10% of their crown showing symptoms can be considered for treatment. Extent of symptoms are an indicator of possibility of successful therapy, but expressions such as percent of crown affected are not necessarily an accurate measure of the distribution of *C. ulmi* in the tree. For example, a tree infected via root graft may have very minimal symptoms, yet it is incurable (GNL observations). Rather than using percent of crown symptomatic, Campana (1) related rate of success to the length of clear sapwood between the cut and the lowest DED-caused streaking on the pruned limb; his success rate (percent of pruned trees that remained free of DED symptoms through the successive two growing seasons) was 87% when 3 m or more of clear wood was obtained and only 12% when this distance was 1.5 m or less. Highest rates of therapeutic success were obtained when a combination of pruning and stem injection of fungicides were administered (3, 4, 5).

In this paper I will relate the results of operational therapeutic treatments of 82 diseased elms, emphasizing accurate assessment of the infection within the tree and the delivery of fungicide directly into the diseased section.

### Materials and Methods

The data presented here represent therapeutic treatments of diseased curb side and garden elms in Syracuse NY (31 trees, 1975-1983) and the Washington DC area (51 trees, 1981-1983). Data on tree parameters, foliar symptoms, sapwood streaking, types, amounts of fungicides injected, and descriptions of pruned limbs were recorded when treatments were made. Results were assessed by inspection (and sometimes sampling) of all trees in late summer or early fall of the year following therapeutic treatment. In six elms given further treatments, the final analyses of their conditions were made in the year following

the last treatment.

Treatments employed included pruning of diseased limbs, injection of benzimidazole fungicides in the root collar region or into diseased areas in the large limbs or the main bole, or combinations of these. Pruned limbs were always debarked to determine the lower limit of sapwood DED-induced streaking. If the distance between the basal cut and the lowest visible streaking was less than 3 m, the limb system was pruned at a lower crotch or the tree was considered for injection. In some cases the DED streak was traced downward by lifting 2-4 cm long V-shaped bark flaps cut with a sharp knife after the outer bark had been removed with an axe. Early injections in Syracuse utilized carbendazim phosphate (similar to LIGNASAN-BLP®), prepared in my laboratory from Benomyl, injected in amounts of active ingredient equivalent to 10-100 times the dosage recommended for the commercial product. We used a commercial formulation of thiabendazole (ARBOTECT-20S®, Merck & Co. Inc., Rahway, NJ) when it became available because it was as active against *C. ulmi* as carbendazim, but labeled for injection at much higher concentrations. Root flare injections of Arbotect were delivered at the label rate of 4 oz (118g) of formulation (26.6% AI, diluted 1:40 in tap water) per 5 inches of tree diameter for trees over 10" (25 cm) in diameter and at about half that rate for smaller trees. Arbotect was injected into limbs at the rate of 0.2-0.4 oz (28-56 g) per inch (2.5 cm) diameter. DED was confirmed by sapwood streaking on all pruned trees and about 50% of the trees that were injected without pruning. In about seven trees with atypical symptoms, the presence of *C. ulmi* was confirmed by culture of sapwood samples on potato dextrose agar. Year of infection was ascertained by dissection sections with the most advanced symptoms to determine the earliest annual ring that had DED streaking and by extrapolation from these dissections to foliar symptoms in other trees.

Fungicide solution was delivered at less than 2 atmospheres of pressure from 20 L gravity (Dewill Inc., Elmhurst IL) or 12 L pressure (Smith & Co., Utica NY) plastic tanks through ¼" holes drilled 1-2 cm into the sapwood. Crown and upper bole injections required climbing and roping of a tank in

the tree.

The techniques used on the various trees were chosen at the time of treatment and based upon evaluation of the symptoms and upon practical limitations of what we able to do and the quantities of fungicide on hand. We did not randomize treatments nor assign some trees to experimental control groups.

## Results

In both Syracuse and Washington, approximately 90% of trees that were given therapy during the year they were infected (= current year infections) and about 50% of treated elms infected in previous years (= residual infections) were free of DED symptoms the year following treatment. Data for the two areas are combined in Table 1.

Pruning without injection was successful in each of 10 current year infections, but in none of 3 residual infections. Pruning alone was applied only when there was a distance of 3 m or more from the last visible streak to the distal cut; failure of pruning residual infections may have resulted from layering of uncolored new sapwood over infected wood. It is clear that fungicide injections is advisable for any tree pruned after the initial year of infection.

Fungicide injection without pruning succeeded in 76% of current year and 33% of residual infections. All 5 of the injected elms with current year infections that eventually died of DED were young (<20 cm dbh) and had infections well established in the bole at the time of treatment. However, in two other such cases, therapeutic injections succeeded.

Fungicide injection plus pruning was successful therapy for 100% of the current year and 71% of the residual infections. In five of the elms with

DED streaks 1-15 cm wide at the last possible pruning cut, fungicide was injected directly into the streaked and the adjacent wood as well as into the root collars of the trees. In this manner we were to cause the injected liquid to "bleed" from the pruning cut, as was advocated by Kondo (7). All of the trees treated in this manner recovered, although one of them required additional pruning and reinjection the following year.

Relatively massive injection of fungicide in roots of trees with root graft-transmitted DED slowed symptom progression but did not cure any of the 4 trees that we treated.

## Discussion

Decisions for the delivery of fungicide injection for DED therapy is best based on whether the tree in question has a current year or a residual infection, rather than on the percentage of the crown with foliar symptoms. The degree of success reported in operational use of benzimidazole fungicides for therapy of Dutch elm disease has been related to the extent of foliar DED symptoms at the time of treatment (8), delivery of an appropriate dosage of active ingredient (10), distribution of compound within the tree (9, 11), and susceptibility of the strain of the DED-causing fungus (*Ceratocystis ulmi*) to the fungicide (5). Manufacturer's guidelines generally state that elms showing 5-10% of the crown symptomatic are suitable candidates of therapeutic injections. While useful, this simplistic approach can lead to disappointment when trees in which there is little possibility of success are injected or in unnecessary condemnation of other trees that might be saved.

The assumption that elms with 5% crown symptoms of DED can be effectively treated is valid if

**Table 1. Results of operational Dutch elm disease therapy: (number) of trees treated and percent without symptoms in the year following last treatment**

Infection type	Treatments						
	Prune		Inject		Prune & inject		Total
Current year	(10)	100	(21)	76	(18)	100	(49) 89.8
Residual	(3)	0	(12)	33	(14)	71	(29) 48.3
Root graft	(0)	-	(4)	0	(0)	-	(4) 0
Total	(13)	77	(37)	54	(32)	88	(82) 70.7

the DED assessed is the result of an elm bark beetle inoculation *during the current year*; it does not pertain to infections that occurred in previous years or to infections that occurred through root grafts. Any DED symptoms that appear within eight weeks of foliation are almost certainly those of residual or root graft infections. Symptoms that first appear in more than one limb system either reflect two or more independent beetle-induced infections, systemic DED, or root graft transmission. Conversely, a systemic or root graft infections sometimes are manifest by DED symptoms in a very limited section of the crown, especially in low vigor trees. For these reasons assessment of DED symptoms is best done by following *C. ulmi*-induced streaking of the sapwood as revealed by debarking pruned branches and/or small cuts through the bark of the intact tree.

Even intensive root and root collar injections may result in sporadic distribution of active ingredient in large and complex elm trees. It seems possible that achievement of sufficient concentrations of fungicide in diseased sections of the tree may be further complicated by a reduction of water movement associated with wilting. In order for root collar injections to deliver a therapeutic dosage of fungicide at the site of the infection high in the tree, it is probably necessary to inject excessive amounts of material of which the vast majority is deposited away from the site of the infection. Delivery of fungicide directly to the infected wood probably greatly increases the rate of therapeutic success. The five trees that we streak-injected would have been considered hopeless cases according to conventional guidelines, yet all remained free of DED symptoms after the treatments were terminated. Since the work summarized here we have apparently cured elms of DED by injecting small amounts (2-10 oz or 56-280 g of formulation) of Arbotect into the streaked wood without injecting at the root collar. On the other hand, our pruning of large elms was usually successful; I agree with Campana's (2) statement that "DED can most easily be eradicated from large trees with small infections".

Contrary to conventional wisdom, residual DED infections can be successfully treated, even if the fungus has reached, to a limited extent, the main bole. Rate of success of therapeutic injections is

considerably enhanced by pruning and, especially, by direct injection of the streaked wood within the crown of the tree. Nevertheless, the most important ingredient for successful DED therapy is detection of new infections when first symptoms appear.

**Acknowledgments.** Technical assistance was provided by Wayne Jones, Michael Griggs, Alan Jones and Alison Teale. Merck & Co. provided, free of charge, some of the fungicide used in this work.

### Literature Cited

1. Campana, R.J. 1976. *Tracing Dutch elm disease infections for depth of infection following excision of infected branches*. Am. Phytopathol. Soc. Proc. 2:95 (Abstr.).
2. Campana, R.J. 1978. *Eradicative pruning*. Pp. 33-34 in Campana, R.J., and Sinclair, W.A. (eds.). *Dutch elm disease: Perspectives after 60 years*. Search (Agriculture) 8(5):52 p.
3. Campana, R.J. and Gregory, G.F. 1976. *Dutch elm disease control by pressureinjected solubilized MBC HCL and excision of infected branch systems*. Proc. Am. Phytopath. Soc. 3:324 (Abstr.).
4. Gregory, G.F. and Allison, J.R. 1979. *The comparative effectiveness of pruning versus pruning plus injection of trunk and/or limb for therapy of Dutch elm disease in American elms*. J. Arboric. 4:1-4.
5. Gregory, G.F. 1982. *Comparative Dutch elm disease therapy: Pruning of symptomatic limbs following Arbotect 20-S and Lignasan BLP injection*. Pp 486-497 in Kondo, E.S., Hiratsuka, Y., Denyer, W.B.G. (eds.) *Proceedings of the Dutch Elm Disease Symposium and Workshop*. Manitoba Natural Resources, Winnipeg.
6. Himelick, E.B. and Cepelcha, D.W. 1976. *Dutch elm disease eradication by pruning*. J. Arboric. 2:81-84.
7. Kondo, E.S. and Huntley, G.D. 1973. *Root-injection field trials of MBC-phosphate in 1972 for Dutch elm disease control*. Can. For. Serv. Inf. Rep. O-X-182. 17 p.
8. Lanier, G.N. 1987. *Fungicides for Dutch elm disease: Comparative evaluation of commercial products*. J. Arboric. 13:189-195.
9. Nishijima, W.T. and Smalley, E.B. 1979. *Distribution and persistence of systemic fungicides in trunk injected elms*. Pp 151-164 in Kielbaso, G. (ed.) *Proc. Symposium on Systemic Chemical Treatment of Trees*. Braun-Brumfield, Ann Arbor. 357 p.
10. Smalley, E.B. 1978. *Systemic chemical treatments of trees for protection and therapy*. Pp 34-35. In Campana, R.J., and Sinclair, W.A. (eds.). *Dutch elm disease: Perspectives after 60 years*. Search (Agriculture) 8(5):52 p.
11. Stennes, M.A., and French, D.W. 1987. *Distribution and retention of thiabendazole hypophosphite and carbendazim phosphate injected into mature American elms*. Phytopath. 77:707-712.

*SUNY College of Environmental Science and Forestry  
Syracuse, New York 13210*