Rocky Mountain Juniper Seed Source Test in Western Kansas, U.S.

Wayne A. Geyer, Keith D. Lynch, and Charles J. Barden

Abstract. Twenty-six sources of Rocky Mountain juniper (*Juniperus scopulorum* Sarg.) were established as a provenance test near Colby, Kansas, U.S. in 1980. Height, diameter, survival, number of stems, crown density, branch angle, and vigor were evaluated with analysis of variance, cluster analysis, simple correlation, and regression analysis techniques. Source differences were found. Total height at 23 years ranged from 3.6 to 5.4 m (11.9 to 17.8 ft), diameter breast height (dbh) ranged from 5.6 to 10.4 cm (2.2 to 4.2 in), and survival from 10% to 100%. Height growth was weakly related to geographic variables, but dbh was related to latitude, longitude, and elevation. Selection of fast-growing sources may begin at 5 years after field planting.

Key Words. Growth characteristics; juniper; *Juniperus scopulorum*; Rocky Mountain juniper; seed sources.

Rocky Mountain juniper (*Juniperus scopulorum* Sarg.) grows on the western edge of the Great Plains, U.S. as essentially a lower-elevation pine tree, on the upper Puget Sound basin, and on Vancouver Island, British Columbia, at or near sea level. It ranges in size from a small tree to a sprawling shrub on dry mountain sites (Harlow and Harrar 1969). Taxonomic studies have been numerous (Fassett 1944; Comer et al. 1982; Adams 1983; Schaefer 1995). Genetic variation of various major traits were determined for 10-year-old plantings, and seed-source selection zones in the Great Plains were identified by Cunningham and King (2000).

This test is part of a larger regional study initiated by the Technical Committee of the Great Plains Agricultural Council in 1973. The objectives were to determine the genetic variation of major traits and to identify the best sources of seed for windbreak applications. This article presents the portion of the study established in Colby, Kansas after 23 years of growth in the field.

MATERIALS AND METHODS

A complete description of the methodology used is found in Van Haverbeke and King (1990). Seed cones were collected from 26 trees in natural stands and windbreaks across the Great Plains (Figure 1; Table 1). Seed trees that had favorable windbreak characteristics (abundant branches, dense foliage) were selected. Seed lots were sown in the USDA Forest Service Bessey Nursery near Halsey, Nebraska, in 1977 and were outplanted at 12 test sites in 1980. This study site is Colby, Kansas (39.38°N and 101.07°W) located in the central geographic region of zones established by Van Haverbeke

(1968). Soil is from the series of Keith loam (fine-silty, mixed, mesic Aridic Haplustolls) and Richfield silty clay loam (fine, montmorillonitic, mesic Aridic Haplustolls). A randomized complete-block with five replications of four-tree, linear plots was established having 20 trees maximum for each source. A total of 520 trees were planted along with 2320 eastern redcedar (*Juniperus virginiana* L.). Spacings in the plantation were 2.4 m (7.9 ft) within rows and 3.6 m (11.9 ft) between rows. A single row was planted on all sides as a border. Complete cultivation was maintained for 5 years. Growth evaluations were made at 5, 10, and 23 growing seasons at this site.

Statistical analyses were performed with SAS software (Statistical Analysis System; SAS Institute Inc., Cary, NC). Data were analyzed by using the analysis of variance procedure and the Duncan's means test was used for mean separation. Cluster analyses (Ward's minimum variance) were used to identify seed sources with similar performance characteristics. Variables used in those analyses were survival, total height, diameter breast height (dbh), number of stems, crown density, branch angle, vigor, latitude, longitude, and elevation. If trees had multiple stems, then only the largest was evaluated for height and diameter. Simple Pearson correlations were computed among seed sources of individual trees for each trait. The correlation of each trait with latitude and longitude was also examined. Least squares regression models of differences in latitude, longitude, and elevation between source locations and the Colby plantation were run to find the extent of geographic area from which seed may be collected to give good tree growth performance in western

Figure 1. Seed selection zones and seed source locations in the northern Great Plains.

Kansas. To determine the reliability of early seed source selection, age/age correlations were computed for survival and height of source means at age 10 and age 23.

RESULTS

Significant differences among seed sources existed for survival, height, dbh, number of stems, crown density, branch angle, and vigor. Each is discussed separately, but we also chose to identify more general patterns of variation by using cluster analysis.

Survival

Seed sources were significantly different (analysis of variance) in survival rates through age 23 with a range of 10% to 100% (Table 2). Total survival across the planting was 72.7%. No specific patterns were indicated geographically, but the best surviving sources (70% to 100%) generally were from Nebraska, North Dakota, and Montana. The poorest source was from Wyoming (10%). No trends for survival were found with latitudes, longitudes, or elevations alone.

Height and Diameter

No specific patterns were indicated geographically for either total height or dbh. The mean height and dbh for all trees were 4.7 m (15.5 ft) and 8.6 cm (3.4 in). For all heights and

diameters, the analysis of variance (Table 2) was significant for differences among sources ($P = 0.01$) with heights ranging from 3.6 to 5.4 m (11.9 to 17.8 ft) and dbh ranging from 5.3 to 12.4 cm (2.1 to 5 in). The tallest trees in the top 20% had high survival (at least 80%) and were in the upper 30% of dbh (9.4 cm [3.8 in]). The largest diameter trees were from the New Mexico Source 8302, but survival was only 40%. The tallest trees had better survival ($\geq 80\%$).

Other Variables

The mean value of the number of stems per tree for the entire planting was 2 with a range of 1.3 to 2.6 stems per tree. Only two sources showed a consistent trend toward a single-stem habit: 3312 and 3313 from northwest Wyoming. Crown density was good for all trees planted. Values ranged from medium to dense for windbreak characteristics. The branch angle was mostly between 45° and 90° . A few sources (3312 and 3313) had a more acute angle. All surviving specimens appeared healthy after 23 years growth even under the drought regimes experienced the last few years in northwestern Kansas.

Table 1. Origin of 26 Rocky Mountain juniper seed sources planted at Colby, Kansas, in 1980.

Source no.	Local area state	Latitude ($^\circ$ N)	Longitude ($^\circ$ W)	Elevation (ft)
3213	Tensleep, WY	44.10	107.50	4100
3312	Muddy Gap, WY	42.35	107.46	4695
3313	Rawlins, WY	41.80	107.22	7220
4611	Big Timber, MT	45.85	109.96	4265
4612	Columbus, MT	45.64	109.27	4100
5311	Sidney, MT	47.73	104.14	1970
5314	Culbertson, MT	48.16	104.52	2050
5511	Towner, ND	48.36	100.41	1475
5513	Towner, ND	48.36	100.41	1475
5611	Larimore, ND	47.92	97.60	1150
5613	Larimore, ND	47.92	97.60	1150
5614	Larimore, ND	47.92	97.60	1150
5811	Mosby, MT	47.00	107.89	2705
5813	Mosby, MT	47.00	107.89	2705
5821	Billings, MT	45.80	108.52	3280
5822	Hathaway, MT	46.28	106.20	2790
5824	Custer, MT	46.13	107.55	2790
5826	Sarpy, MT	46.30	107.11	3935
5846	Reva, SD	45.56	103.10	3115
5847	Reva, SD	45.56	103.10	3115
5851	Johnson City, WY	44.06	106.59	4590
6411	Chadron, NE	42.85	102.99	3445
6412	Chadron, NE	42.85	102.99	3445
6413	Chadron, NE	42.85	102.99	3445
6712	Casebier, WY	42.33	104.51	5085
8302	Ruidoso Downs, NM	33.35	105.68	7380

Table 2. Survival, height, and dbh rankings of seed sources at 23 years.

Source no.	Survival (%)	Height m (ft)	Range m (ft)	dbh cm (in)	Range cm (in)
5826	80 bcd^z	5.4 (17.7) a^z	4.3–6.4 (14.0–21.0)	10.4 (4.1) bc^z	2.5–15.2 (1.0–6.0)
6411	100 d	5.3 (17.4) ab	4.3–6.1 (14.0–20.0)	9.7 (3.8) bcd	7.6–12.7 (3.0–5.0)
6412	90 cd	5.3 (17.3) ab	4.4–6.6 (14.5–21.5)	10.4 (4.1) bc	5.1–17.8 (2.0–7.0)
5311	80 bcd	5.3 (17.3) abc	4.0–6.4 (13.0–21.0)	10.4 (4.1) b	5.1–17.8 (2.0–7.0)
5846	90 cd	5.2 (17.0) abcd	4.7–5.6 (15.5–18.5)	9.4 (3.7) bcde	5.1–12.7 (2.0–5.0)
5513	40 abc	5.2 (16.9) abcd	4.1–5.8 (13.5–19.0)	10.2 (4.0) bc	5.1–12.7 (2.0–5.0)
8302	40 bcd	5.1 (16.6) abcd	4.6–5.8 (15.0–18.9)	12.4 (4.9) a	7.6–15.2 (3.0–6.0)
6712	10 a	5.0 (16.5) abcde	4.1–6.6 (13.5–21.5)	9.9 (3.9) bc	5.1–15.2 (2.0–6.0)
5813	90 cd	5.0 (16.5) abcdef	4.4–5.6 (14.5–18.5)	8.9 (3.5) bcde	5.1–12.7 (2.0–5.0)
5613	90 cd	5.0 (16.4) abcdef	4.6–5.3 (15.0–17.5)	9.1 (3.6) bcde	5.1–12.7 (2.0–5.0)
3312	80 bcd	5.0 (16.3) abcdef	4.1–5.5 (13.5–18.0)	9.4 (3.7) bcde	5.1–12.7 (2.0–5.0)
5614	80 bcd	4.9 (16.0) abcdefg	4.3–6.1 (14.0–20.0)	9.1 (3.6) bcde	5.1–12.7 (2.0–5.0)
6413	90 cd	4.9 (16.0) bdefg	4.3–5.9 (14.0–19.5)	8.6 (3.4) cdef	5.1–12.7 (2.0–5.0)
5611	90 cd	4.8 (15.9) bcdefg	4.3–5.3 (14.0–17.5)	9.7 (3.8) bcd	5.1–15.2 (2.0–6.0)
5847	40 bcd	4.8 (15.9) bcdefg	4.4–5.3 (14.5–17.5)	7.9 (3.1) def	5.1–10.2 (2.0–4.0)
5821	100 d	4.8 (15.6) cdefgh	4.0–5.2 (13.0–17.0)	8.1 (3.2) def	5.1–10.2 (2.0–4.0)
5824	80 bcd	4.8 (15.6) defgh	3.8–5.5 (12.5–18.0)	7.6 (3.0) efg	5.1–12.7 (2.0–5.0)
5822	70 bcd	4.7 (15.5) defgh	4.1–5.6 (13.5–18.5)	8.1 (3.2) def	5.1–10.2 (2.0–4.0)
5511	40 bcd	4.7 (15.4) defgh	4.0–5.6 (13.0–18.5)	8.6 (3.4) bcde	5.1–15.2 (2.0–6.0)
3313	75 bcd	4.5 (14.8) efgh	4.0–5.2 (13.0–17.0)	7.6 (3.0) efg	5.1–10.2 (2.0–4.0)
5314	40 bcd	4.5 (14.8) fgh	3.7–5.0 (12.0–16.5)	8.6 (3.4) bcde	7.6–12.7 (3.0–5.0)
5811	30 ab	4.5 (14.6) gh	3.4–5.2 (11.0–17.0)	8.1 (3.2) def	5.1–10.2 (2.0–4.0)
3213	85 cd	4.3 (14.2) h	3.7–5.5 (12.0–18.0)	6.9 (2.7) fgh	2.5–10.2 (1.0–4.0)
5851	40 abc	3.8 (12.6) i	3.0–4.4 (10.0–14.5)	5.8 (2.3) h	2.5–7.6 (1.0–3.0)
4611	70 bcd	3.8 (12.5) i	2.9–4.3 (9.5–14.0)	6.1 (2.4) gh	5.1–10.2 (2.0–4.0)
4612	90 cd	3.6 (11.8) i	3.2–4.6 (10.5–15.0)	5.6 (2.2) h	2.5–7.6 (1.0–3.0)

^zValues followed by the same letters within a column are not significantly different at the $P = 0.05$. **Bold type indicates tallest 20%.**

Cluster Analysis

Cluster analysis included all 26 sources. The analysis differentiated two, three, and four clusters with significance that was similar for survival, height, and dbh. The other variables listed were not significant and were therefore deleted in the analysis. General patterns of variation were identified by using this analysis. No distinct geographic patterns were shown in the three- and four-cluster combinations, so the two-cluster analysis was used. The height values for first and second clusters were 3.2 and 4.9 m (10.6 and 16.2 ft), dbh values were 6.3 and 9.9 cm (2.5 and 4 in), and survival rates were 66.7% and 73.9%, respectively. Sources 4611, 4612, and 5851 were the shortest. These were collected from southwestern Montana and north central Wyoming. In addition, all three sources were also separated from the others in the three and four combinations.

Correlations

Age/age correlations between height and dbh were particularly high. Height growth at 5 years predicted the tallest trees at 10 years (Pearson correlation coefficient = 0.960). Height growth at 10 years (Table 3) predicted growth at 23 years

(0.895). Five-year data were not available for determining 23-year relationships, but one could assume that the correlations would be high. The tallest trees had the largest diameters (correlation = 0.858). The other major variables and the geographic traits did not provide any significantly high correlations to predict growth (Table 3).

Regression Analysis

The reliability of the regression model to predict the 23-year height and diameter from the 10-year data by using a com-

Table 3. Correlations of height and dbh of Rocky Mountain juniper with age and geographic traits.

	Height at 10 yr	Height at 23 yr	dbh at 23 yr	Long.	Lat.	Elev.
Height 5 yr	0.960 ^z					
10 yr		0.895	0.909	0.267	-0.144	0.175
23 yr			0.858	0.332	-0.097	-0.164
dbh 23 yr		0.858		0.300	-0.296	-0.050

^zNote: Approximate value from Cunningham and King (2000).

bination of the geographic variables gave R-square values of 0.3755 (height) and 0.6721 (dbh) for values significant at the 1% level. Thus, the height model was weak, but the dbh model was relatively strong. Only longitude squared was significant in the height prediction. Latitude, longitude, elevation, and the squares of longitude and elevation were significant in predicting dbh. Northerly latitudes, easterly longitudes, and elevations lower than those at the Colby site produced larger diameter trees. All but one source came from locations north of Colby.

DISCUSSION AND CONCLUSIONS

In an earlier study in the Great Plains region (Cunningham and King 2000), Rocky Mountain juniper trees survived better, were shorter, and had smaller crowns than did eastern redcedar (*J. virginiana* L.) at 10 years of age. Rocky Mountain juniper has better crown compactness for wind reduction than redcedar. In our Kansas study, we found, at 23 years of age that only three seed sources (4611, 4612, and 5851) had a geographic difference from the others sources compared. They were from southwestern Montana and north central Wyoming and were much shorter in height. These sources are not recommended for windbreak plantings in western Kansas. Thus, the general good health, dense crown, and many stems of Rocky Mountain juniper would provide excellent structure, and this species should be selected for use in establishing new windbreaks in western Kansas and nearby areas in eastern and western Nebraska. Our findings showed growth at 10 years predicted superior growth at 23 years and in agreement with Schaefer (1995) and VanHaverbeke and King (1990) for 5-year and 10-year relationships. Rocky Mountain juniper is prone to *Cercospora* needle blight in more humid areas of the eastern Great Plains and is not recommended for windbreak plantings in this area.

Acknowledgments. This is Contribution no. 07-69-J from the Kansas Agricultural Experiment Station, Manhattan Kansas.

LITERATURE CITED

- Adams, R.P. 1983. Intraspecific terpenoid variation in *Juniperus scopulorum*: Evidence for Pleistocene refugia and recolonization in western North America. *Taxon* 32: 30–46.
- Comer, C., R.P. Adams, and D.F. Van Haverbeke. 1982. Intra- and interspecific variation of *Juniperus virginiana* and *J. scopulorum* seedlings based on volatile oil composition. *Biochemical Systematics and Ecology* 10:179–306.
- Cunningham, R.A., and R.M. King. 2000. Juniper seed sources in the Great Plains. General Technical Report, RMRS-GTR-51. USDA Forest Service, Rocky Mountain Forest Experiment Station. Fort Collins, CO.
- Fassett, N.C. 1944. *Juniperus virginiana*, *J. horizontalis* and *J. scopulorum*—II. Hybrid swarms of *Juniperus virginiana* and *J. scopulorum*. *Bulletin of the Torrey Botanical Club* 71: 475–483.
- Harlow, W.M., and E.S. Harrar. 1969. *Textbook of Dendrology*, 5th Edition, McGraw Hill Book Co., Inc., New York. 542 pp.
- Schaefer, P. 1995. Ten-year results of an eastern redcedar and Rocky Mountain juniper provenance test in eastern South Dakota. *Northern Journal of Applied Forestry* 12:30–35.
- Van Haverbeke, D.F. 1968. A population analysis of *Juniperus* in the Missouri river Basin. New Series 38. Lincoln, NE, University of Nebraska Studies. 82 pp.
- Van Haverbeke, D.F., and R.M. King. 1990. Genetic variation in the Great Plains *Juniperus*. USDA Forest Service. Res. Pap. M-292. Lincoln, NE. 8 pp.

Wayne A. Geyer (corresponding author)
Professor
Forestry Division
Throckmorton Hall
Kansas State University
Manhattan KS 66506, U.S.
wgeyer@ksu.edu

Keith D. Lynch
Associate Professor
Forestry Division
Throckmorton Hall
Kansas State University
Manhattan KS 66506, U.S.

Charles J. Barden
Associate Professor
Forestry Division
Throckmorton Hall
Kansas State University
Manhattan KS 66506, U.S.

Résumé. Trente-six provenances de genévrier des Rocheuses (*Juniperus scopulorum* Sarg.) ont été plantées dans le cadre d'un test de provenance près de Kolby au Kansas en 1980. La hauteur, le diamètre, le taux de survie, le nombre de tiges, la densité de la cime, l'angle des branches et la vigueur ont été évalués au moyen de techniques d'analyses de variances, d'analyse de grappes, de corrélations simples et d'analyses de régression. Des différences entre les provenances ont été trouvées. La hauteur totale après 23 ans se situait entre 3,6 et 5,4 m, le DHP entre 5,6 et 10,4 cm, et le taux de survie entre 10 et 100%. La croissance en hauteur était faiblement corrélée aux variables géographiques, mais le DHP était relié à la latitude, la longitude et l'élévation. La sélection de provenances à croissance rapide pourrait se faire à compter de la cinquième année après la plantation en plein champs.

Zusammenfassung. Im Jahr 1980 wurden in Colby, Kansas, 26 Quellen für einen Herkunftstest von Wacholder (*Juniperus scopulorum* Sarg.) eingerichtet. Mit ANOVA wurde die Höhe, Durchmesser, Anzahl der Stämme, Astwinkel, Vitalität getestet, eine Cluster-Analyse, simple Korrelation und Regressionsanalysen-Technik. Es wurden Differenzen zwischen den Quellen gefunden. Die totale Höhe im 23. Jahr reichte von 3,6 bis zu 5,4 m, der Brusthöhendurchmesser reichte von 5,6 bis 10,4 cm und die Überlebensrate von 10 bis 100 %. Das Höhenwachstum war schwach mit der geografischen Lage verbunden, aber der Brusthöhendurchmesser stand in Beziehung zu Länge, Breite und Höhe. Eine Auswahl der schnellwachsenden Quellen kann ab dem 5. Wachstumsjahr erfolgen.

Resumen. En 1980, se establecieron 26 fuentes de junípero Rocky Mountain (*Juniperus scopulorum* Sarg.) en una prueba cerca de Colby, Kansas. Se evaluó altura, diámetro, supervivencia, número de tallos, densidad de copa, ángulo de ramas y vigor con técnicas ANOVA, análisis cluster, correlación simple y análisis de regresión. Se encontraron diferencias entre las fuentes. La altura total a 23 años fue de 3.6 a 5.4 m (11.8 a 17.7 pies), el dap (diámetro a la altura del pecho) varió de 5.6 a 10.4 cm. (2.2 a 4.1 pulgadas), y la supervivencia de 10 a 100%. El crecimiento en altura estuvo débilmente relacionado a las variables geográficas, pero el dap estuvo relacionado con latitud, longitud y elevación. La selección de fuentes de rápido crecimiento puede empezar a los cinco años después de la plantación en el campo.