Technical Note**TREES AND WIND: WIND SCALES AND SPEEDS****by Scott Cullen**

Key Words. Trees and wind; wind; wind scales; wind speeds.

NEED AND PURPOSE

Arborists and urban foresters frequently refer to “wind loads,” “sail areas,” and so forth when discussing the risk of tree failure. Whether the discussion is generally descriptive or specifically quantitative, wind speed is an essential consideration. Wind load is a function of wind speed. Acceptable levels of risk must be associated with wind speeds and their probability of recurrence (Cullen 2002a).

It is conventional in scholarly literature to cite wind speed in SI (*Système International*) units of m/s (NIST, no date). It is customary, however, for meteorological agencies to report wind speeds in miles per hour, knots, or kilometers per hour. The preferred units vary by both agency and country. In addition, various classification scales are used to describe weather events by wind speed ranges. Some of the tree-risk literature employs these scale numbers (e.g., Sinn and Wessolly 1989; Wessolly 1995; Peterson and Rebertus 1997; Wessolly and Erb 1998; Hayes 1999; Peterson 2000; Brudi 2002).

These various usages may be an obstacle to proper understanding of the tree-wind literature (Cullen 2002b), to practical tree-risk management, and to dissemination of knowledge and methods across national and cultural boundaries. The tables in this article should facilitate simple and accurate comparison.

COMPARING SCALES AND SPEEDS

A comparison of wind speed scales and wind speeds in various units of measure is presented in Table 1. The table is not intended to catalog all wind speed scales, but includes several common systems. The World Meteorological Organization provides a list of weather reporting agencies around the world (WMO, no date) that employ various systems.

Some scales incorporate other elements of storm severity in addition to wind speed. For example, the Fujita Scale (Fujita 1987; NCDC 2001a; TTU 2002)—or Fujita-Pearson Scale—(Stormfax 2001) considers the width and length of tornado damage paths; and the

Dolan-Davis Northeaster Storm Scale (Dolan and Davis 1992) considers the “storm surge” in coastal water bodies. A variety of scales are catalogued by the Natural Hazards Research Centre (2001). These scales may be useful in assessing risks of tree damage in particular locales.

TREES AS WIND SPEED INDICATORS

A number of scales have been developed to use tree movement or deformation as an indication of current or experienced wind speed.

Tree Movement and Damage

Historically, the Beaufort Scale was developed to scale wind speeds over water and describe wind effects that could be observed by mariners. The scale has been adjusted for use on land, including description of wind effects on trees. The Fujita Tornado Scale and Saffir-Simpson Hurricane Scale (NWS, no date-b; Simpson and Riehl 1981; NCDC 2001b) use tree damage as an index of wind speeds. These effects are presented in Table 2. Rating damage using the Fujita Scale has been characterized as “highly subjective and variable” (Edwards and Harmon, no date; Doswell and Burgess 1988).

Deformation of Trees

In chronically windy locations, wind can also affect tree growth resulting in permanent deformation. Robertson (1987) notes that a number of investigators have developed indices of deformation related to wind speeds. These indices include the Griggs-Putnam index (Putnam 1948) for North American conifers; the Barsch index (Barsch 1963) for European broadleaves; and Yoshino (1975) and Yoshino et al. (1976) for European and Japanese trees. Koepl (1982) reports that the Griggs-Putnam index was further developed to scale deformation against wind characteristics including mean annual wind speed, mean growing season wind speed, mean nongrowing season wind speed, and percentage of winds from prevailing direction. The predictive reliability of these indices is not without critics (Hennessey 1980).

Other studies of tree deformation by chronic wind include those by Cordero (1999), Musselman et al. (1990), and Noguchi (1979, 1992).

Table 1. Wind scales and speeds (copyright Scott Cullen 2002).

Scale				
Fujita Tornado ¹	Saffir / Simpson		Tropical Cyclone	
	Atlantic ² NE. Pacific	NW Pacific Typhoon ³ Indian Cyclone (MSW) ⁴	AUS ⁵ km/h	Hong Kong ⁶ km/h
Description	Category	Hurricane		
<div>$V_{WIND} = 14.1 \bullet (F + 2)^{1.5}$</div>				
Gale Tornado	F0			Tropical Depression
				Tropical Storm
Moderate Tornado	F1			Severe Tropical Storm
				Typhoon
Significant Tornado	F2	1	1 Minimal Typhoon	
		2	2 Moderate Typhoon	
		3	3 Strong Typhoon	
		4	4 Very Strong Typhoon Super Typhoon	
Severe Tornado	F3	5	5 Devastating Typhoon	
Devastating Tornado	F4			
Incredible Tornado	F5			
None Expected	F6....			
	F12			
	Mach 1			

¹(NCDC 2001a). ²(Landsea 2000a; NWS, no date-a). ³(Landsea 2000c; Navy, no date; NWS 1999). ⁴MSW classified by higher peak gusts (Navy-no date). ⁵(BoM-AU 2001); strongest gust. ⁶(Hong Kong Observatory near center over 10 minutes. ⁷(Hong Kong Observatory 1999a; Landsea 2000a; NWS, no date-b) NWS-NCDC 2000; NIST, no date). ⁹Calculated speeds > 4 m/s rounded to nearest whole number.

Scale			Wind Speed			
Beaufort ⁷			(Published Calculated ⁸)			
Force Number	Description					
	(US-NWS)	(WMO)	MPH	Knots	km/h	m/s ⁹
0	Calm	Calm	0 - 1	< 1	< 1	0 - 0.4
1	Light	Light Air	1 - 3	1 - 3	1 - 6	0.4 - 1.3
2	Light	Light Breeze	4 - 7	4 - 6	7 - 12	1.8 - 3.1
3	Gentle	Gentle Breeze	8 - 12	7 - 10	13 - 19	4 - 5
4	Moderate	Moderate Breeze	13 - 18	11 - 16	20 - 30	6 - 8
5	Fresh	Fresh Breeze	19 - 24	17 - 21	31 - 39	8 - 11
6	Strong	Strong Breeze	25 - 31	22 - 27	40 - 50	11 - 14
7	Strong	Near Gale	32 - 38	28 - 33	51 - 62	14 - 17
			< 39	< 34	< 63	< 17
			39 - 49	26 - 43	63 - 79	17 - 22
8	Gale	Gale	39 - 46	34 - 40	63 - 74	17 - 21
			40 - 72	35 - 62	64 - 116	18 - 32
9	Gale	Strong Gale	47 - 54	41 - 47	75 - 87	21 - 24
10	Whole Gale	Storm	55 - 63	48 - 55	88 - 102	25 - 28
11	Whole Gale	Violent Storm	64 - 72	56 - 63	103 - 117	29 - 32
			50 - 73	44 - 63	80 - 117	22 - 33
12	Hurricane	Hurricane	73 <	64 <	118 <	33 <
			73 - 112	63 - 97	117 - 180	33 - 50
			74 - 95	64 - 82	119 - 153	33 - 42
			< 78	< 67	< 125	< 35
			78 - 106	67 - 92	125 - 170	35 - 47
			96 - 110	83 - 95	154 - 177	43 - 49
			106 - 140	92 - 121	170 - 225	47 - 63
			< 112	< 97	< 180	< 50
			111 - 130	96 - 113	179 - 209	50 - 58
			113 - 157	98 - 136	182 - 253	51 - 70
			131 - 155	114 - 135	211 - 249	59 - 69
			150 <	130 <	241 <	65 <
			140 - 175	121 - 151	225 - 280	63 - 78
			156 <	136 <	251 <	70 <
			156 - 194	136 - 170	251 - 312	70 - 87
			158 - 206	137 - 179	254 - 331	71 - 92
			175 <	151 <	280 <	78 <
			< 206	< 178	< 331	< 92
			207 - 260	180 - 226	333 - 418	93 - 116
			261 - 318	227 - 276	420 - 512	117 - 142
			319 <	277 <	513 <	143 <
			< 738	< 638	< 1187	< 330
			738 <	638 <	1187 <	330 <

=maximum sustained wind for 1 minute (Landsea 2000c); these typhoon categories are also
 atory 1999b) following World Meteorology Organization scheme; average sustained speed in km/h
 = U.S. National Weather Service; WMO=World Meteorological Organization. ⁸(Landsea 2000b;

Table 2. Wind speeds and trees (compiled by Scott Cullen 2002).

Scale			Wind Speed MPH ⁴
Fujita Tornado ¹		Beaufort ²	
Description	Category	Force Number	
Tree Effect		Tree Effect	
		0	0 - 1
		1	1 - 3
		2	4 - 7
		3	8 - 12
		4	13 - 18
		5	19 - 24
		6	25 - 31
Gale Tornado	F0	7	32 - 38
		8	39 - 46
		9	47 - 54
		10	55 - 63
Moderate Tornado	F1	11	64 - 72
		12	73 <
		Saffir / Simpson ³	
		1	74 - 95
		2	96 - 110
Significant Tornado	F2		< 112
			113 <
Severe Tornado	F3		< 157
			158 <
Devastating Tornado	F4		< 206
			207 <
Incredible Tornado	F5		< 260
			261 <
			< 318

¹(NCDC 2001a; Stormfax 2001). ²(Landsea 2000a; NWS, no date-b). ³(Landsea 2000a; NWS, no date-a). ⁴See Table 1 to convert mph to other units.

These indices and studies may be useful in assessing wind exposure of trees and biomechanical effects of winds of varying speeds.

LITERATURE CITED

- Barsch, Dietrich. 1963. Wind, baumform und landschaft / Dietrich Barsch, in Studien zum problem der deformation von baumkronen durch wind. Freiburger Geographische Hefte H1 (Albert-Ludwigs-Universitaet, Geograph. Institut 1, Freiburg).
- Bureau of Meteorology—Australia (BoM-AU). 2001. Cyclone Severity Categories. <http://www.bom.gov.au/info/cyclone/#CycloneSeverityCategories> (accessed 7/24/02).
- Brudi, Erk. 2002. Trees and statics: An introduction. In Smiley, E. Thomas, and Kim D. Coder (Eds.). Proceedings of the Tree Biomechanics Conference, Savannah, GA. International Society of Arboriculture, Champaign, IL (in press).
- Cordero, Roberto A. 1999. Ecophysiology of *Cecropia schreberiana* saplings in two wind regimes in an elfin cloud forest: growth, gas exchange, architecture and stem biomechanics. *Tree Physiol.* 19(3):153–163.
- Cullen, Scott. 2002a. Guying a large tree: Decision making and design (the Bedford protocol). In Smiley, E. Thomas, and Kim D. Coder (Eds.). Proceedings of the Tree Biomechanics Conference, Savannah, GA. International Society of Arboriculture, Champaign, IL (in press).
- Cullen, Scott. 2002b. Trees and wind: A bibliography for tree care professionals. *J. Arboric.* 28(1):41–51. <http://joa.isa-arbor.com/request.asp?JournalID=1&ArticleID=25&Type=1> (accessed 7/24/02).
- Dolan, R., and R.E. Davis. 1992. Rating northeasters. *Mariners Weather Log* 36(1):4–11. (Cited in Hilton Head Flood Hazard Mitigation Plan. 1995. Town of Hilton Head, Hilton Head, SC. <http://www.ci.hilton-head-island.sc.us/projects/flood/e.html> (accessed 7/24/02).
- Doswell, C.A., and D.W. Burgess. 1988. On some issues of United States tornado climatology. *Monthly Weather Review* 116:495–501.
- Edwards, Roger, and D. Greg Harmon. No date. Lubbock F-Scale Exercise on Spencer, SD Tornado Damage. Storm Prediction Center, NOAA, Norman, OK. http://www.spc.noaa.gov/misc/spencer/spenf_ex.htm (accessed 7/31/02).
- Fujita, T. Theodore. 1987. U.S. Tornadoes, Part 1: 70-Year Statistics. Satellite and Mesometeorology Research Project (SMRP), Research Paper Number 218. University of Chicago. 122 pp.
- Hayes, Ed. 1999. Patterns of tree failure. *Tree Care Indus.* X(4):37–42.
- Hennessey, J.P. 1980. A critique of “trees as a local climatic indicator.” *J. Appl. Meteorol.* 19:1020–1023.
- Hong Kong Observatory. 1999a. Beaufort Wind Scale. <http://www.weather.gov.hk/education/article/009beaufort.htm> (accessed 7/25/02).
- Hong Kong Observatory. 1999b. Classification of Tropical Cyclones. <http://www.weather.gov.hk/informtc/class.htm> (accessed 7/52/02).
- Koeppel, Gerald W. 1982. Wind characteristics and turbine siting, Chapter 2 in Part 2. In Putnam's Power from the Wind (2nd ed.). Van Nostrand Reinhold, New York, NY. 470 pp.
- Landsea, Christopher W. 2000a. How are Atlantic hurricanes ranked? Atlantic Oceanographic and Meteorological Laboratory, NOAA, Miami, FL. (Citing Simpson, R.H., and H. Riehl. 1981. The Hurricane and Its Impact. Louisiana State University Press, Baton Rouge, LA. 398 pp.) <http://www.aoml.noaa.gov/hrd/tcfaq/tcfaqD.html#D1> (accessed 7/25/02).
- Landsea, Christopher W. 2000b. How do I convert from mph to knots and m/s? Atlantic Oceanographic and Meteorological Laboratory, NOAA, Miami, FL. <http://www.aoml.noaa.gov/hrd/tcfaq/tcfaqD.html#D4> (accessed 7/25/02).
- Landsea, Christopher W. 2000c. What is a super typhoon? Atlantic Oceanographic and Meteorological Laboratory, NOAA, Miami, FL. (Citing U.S. Joint Typhoon Warning Center, Guam). <http://www.aoml.noaa.gov/hrd/tcfaq/tcfaqA.html#A3> (accessed 7/25/02).
- Musselman, R.C., G.L. Wooldridge, D.G. Fox, and B.H. Connell. 1990. Using wind-deformed conifers to measure wind patterns in alpine transition at GLEES, pp 80–85. In Schmidt, W.C., and K.J. McDonald (Compilers). Proceedings—Symposium on Whitebark Pine Ecosystems: Ecology and Management of a High-Mountain Resource, Bozeman, Montana, March 29–31, 1989. Intermountain Station General Technical Report INT-270.
- National Institute of Science and Technology (NIST). No date. Guide for the Use of the International System of Units (SI)—Velocity. <http://physics.nist.gov/Pubs/SP811/appenB8.html> (accessed 7/25/02).
- National Climate Data Center (NCDC). 2000. Knots to Miles per Hour Conversion Chart. NOAA, Asheville, NC. <http://www.ncdc.noaa.gov/ol/climate/conversion/windchart.html> (accessed 7/25/02).
- National Climate Data Center (NCDC). 2001a. The Fujita Tornado Scale. NOAA, Asheville, NC. <http://www.ncdc.noaa.gov/ol/satellite/satelliteseye/educational/fujita.html> (accessed 7/25/02).
- National Climate Data Center (NCDC). 2001b. The Saffir-Simpson Hurricane Scale. NOAA, Asheville, NC. <http://lwf.ncdc.noaa.gov/oa/satellite/satelliteseye/educational/saffir.html> (accessed 7/25/02).
- National Weather Service (NWS). 1999. Modified Saffir-Simpson hurricane scale (SSHS) for the northwest Pacific (Appendix A, NWS Operations Manual W/OM12 8/19/99), NOAA, Silver Spring, MD. <http://www.nws.noaa.gov/wsom/manual/archives/NC419903.HTML> (accessed 7/25/02).
- National Weather Service (NWS). No date—a. Saffir-Simpson Hurricane Scale, NOAA, Silver Spring, MD. <http://www.nws.noaa.gov/er/box/Saffir.html> (accessed 7/25/02).

- National Weather Service (NWS). No date–b. Beaufort Wind Scale. Storm Prediction Center, NOAA, Norman, OK. <http://www.spc.noaa.gov/faq/tornado/beaufort.html> (accessed 7/25/02).
- Natural Hazards Research Centre. 2001. Damage Scales. Division of Environmental & Life Sciences, Macquarie University, Sydney, NSW, Australia. <http://www.es.mq.edu.au/NHRC/web/scales/scalesindex.htm> (accessed 7/25/02).
- Navy Guam Weather Watch. No Date. Tropical Cyclones. <http://www.guam.navy.mil/weather.htm> (accessed 7/25/02).
- Noguchi, Y. 1992. Vegetation asymmetry in Hawaii under the trade wind regime. *J. Veg. Sci.* 3(2):223 ff.
- Noguchi, Y. 1979. Deformation of trees in Hawaii and its relation to wind. *J. Ecol.* 67:611–628.
- Peterson, C.J. 2000. Damage and recovery of tree species after two different tornadoes in the same old growth forest: A comparison of infrequent wind disturbances. *For. Ecol. Manage.* 135(1–3):237–252.
- Peterson, C.J., and A. J. Rebertus. 1997. Tornado damage and initial recovery in three adjacent, lowland temperate forests in Missouri. *J. Veg. Sci.* 8(4):559–564.
- Putnam, Palmer Cosslett. 1948. *Power from the Wind*. Van Nostrand-Reinhold, New York, NY. 223 pp.
- Robertson, Alexander. 1987. The use of trees to study wind. *Arboric. J.* 11:127–143.
- Simpson, R.H., and H. Riehl. 1981. *The Hurricane and Its Impact*. Louisiana State University Press, Baton Rouge, LA. 398 pp.
- Sinn, Günter, and Lothar Wessolly. 1989. A contribution to the proper assessment of the strength and stability of trees. *Arboric. J.* 13(1):45–65.
- Stormfax. 2001. Fujita-Pearson Tornado Scale. Stormfax® Weather Almanac. <http://www.stormfax.com/fujita.htm> (accessed 7/25/02).
- Texas Tech University (TTU). 2002. Fujita Tornado Damage Scale. Wind Engineering Research Center, Texas Tech University, Lubbock, TX. <http://www.spc.noaa.gov/faq/tornado/f-scale.html> (accessed 7/25/02).
- Wessolly, Lothar. 1995. Fracture diagnosis of trees, part 3: Boring is no way for reliable fracture diagnosis (including literature citations for parts 1, 2, and 3). *Stadt und Grün* 9:635–640.
- Wessolly, Lothar, and M. Erb. 1998. *Handbüch der Baumstatik und Baumkontrolle* (in German). Patzer-Verlag, Berlin, Germany. 270 pp.
- World Meteorological Organization (WMO). No date. National Weather Services Affected by Tropical Cyclones. WMO, Geneva. <http://www.wmo.ch/web/www/TCP/National%20Warnings.htm> (accessed 7/25/02).
- Yoshino, M.M., M.T. Yoshino, M. Yoshimura, K. Mitsui, K. Urushibara, S. Ueda, M. Owada, and K. Nakamura. 1976. Bora regions as revealed by wind-shaped trees on the Adriatic Coast, pp 59–71. In Yoshino, M.M. (Ed.). *Local Wind Bora*. University of Tokyo Press, Tokyo, Japan.
- Yoshino, M.M. 1975. Wind-shaped trees, Chapter 6.1.3, pp 445–458. In *Climate in a Small Area*. University of Tokyo Press, Tokyo, Japan.

Consulting Arborist
P.O. Box 31152
Greenwich, CT 06831, U.S.
dscottcul@att.net